



ACADIANA GASTROENTEROLOGY

A S S O C I A T E S

439 Heymann Blvd., Lafayette, LA 70503
Ph. (337) 269-0963 * Fax (337) 269-0553
www.acadianagastro.com

- Stephen M. Person, M.D.
- James C. Bienvenu, M.D.
- Erick A. Salvatierra, M.D.
- Christopher P. Herrington, M.D.

- J. Patrick Herrington, M.D.
- Richard K. Broussard, M.D.
- Patrick A. Laperouse, M.D.

MoviPrep BOWEL PREP INSTRUCTIONS FOR YOUR PROCEDURE

BEFORE YOUR EXAM – Purchase the following:

- Prescription for MoviPrep –Your Rx will be sent to the pharmacy 1 week prior to your procedure.
- 4 Gas-X tablets

THE DAY BEFORE YOUR PROCEDURE:

Begin a CLEAR LIQUID DIET when you awaken on Day: _____, Date:_____.

NO SOLID FOODS are to be eaten on this day. You may ONLY HAVE CLEAR LIQUIDS.

Helpful hints to make your prep more tolerable: Refrigerate solution of Moviprep until very cold. Mix Moviprep with up to 2oz. of lemon or lime juice along with ice cold water, add a lid to your cup & drink the solution through a straw.

STEP 1 (Mix First Dose)-

1. Empty 1 Pouch A and 1 Pouch B into the disposable container.
2. Add warm drinking water to the top line of the container and mix to dissolve. Refrigerate.

STEP 2 (Drink First Dose)

1. The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz.), until the full liter is consumed.
2. Drink 16 oz. of a clear liquid of your choice. This is a necessary step to ensure adequate hydration and an effective prep.
3. Follow with 2 (two) Gas-X tablets.

DO NOT EAT OR DRINK ANYTHING after MIDNIGHT, UNLESS INSTRUCTED OTHERWISE, EXCEPT FOR THE PREP THE MORNING OF THE PROCEDURE & A SIP OF WATER WITH BLOOD PRESSURE MEDICATIONS

THE MORNING OF YOUR PROCEDURE, beginning at (time):_____

(If the time of your procedure changes, you must still begin your 2nd dose of prep solution 4 hours before the time of your procedure and fully consume it within 1 hour & 30 minutes)

STEP 3 (Mix Second Dose)

1. Empty 1 Pouch A and 1 Pouch B into the disposable container.
2. Add warm drinking water to the top line of the container and mix to dissolve. Refrigerate.

STEP 4 (Drink Second Dose)

1. The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz.), until the full liter is consumed.
2. Drink 16 oz. of a clear liquid of your choice. This is a necessary step to ensure adequate hydration and an effective prep.
3. Follow with 2 (two) Gas-X tablets.

This entire process is required and is important to follow to ensure a clean colon. If not, your physician will not be able to complete a thorough exam and you may be asked to repeat your procedure.

IF YOU HAVE ANY QUESTIONS ABOUT THE PREP INSTRUCTIONS ABOVE, or
YOUR SCHEDULED PROCEDURE, PLEASE CALL OUR OFFICE: 337-269-0963

www.acadianagastro.com